



69th DAE SYMPOSIUM ON NUCLEAR PHYSICS

DR. B R AMBEDKAR NATIONAL INSTITUTE OF TECHNOLOGY JALANDHAR, PUNJAB



VOLUME 69 (2025)

DECEMBER 8 - 12, 2025

EDITORS

H. KUMAWAT, VINEET KUMAR, P. SHUKLA & ARADHANA SHRIVASTAVA



Sponsored by

Board of Research in Nuclear Sciences,
Department of Atomic Energy, Government of India



Published by:

Scientific Information Resource Division,
Bhabha Atomic Research Centre,
Trombay, Mumbai,
Maharashtra - 400085, India.
Email: headsird@barc.gov.in

Printed by:

Brandskool,
E.D. 119, Dhan Mohalla,
Jalandhar City - 144001 (Punjab) INDIA
Mob : +91-8837600299

69th DAE SYMPOSIUM ON NUCLEAR PHYSICS



**Dr. B. R. Ambedkar National Institute of Technology
Jalandhar, Punjab - 144008**

Volume 69 (2025)

Full proceedings available at: <http://sympnp.org/proceedings/69>

December 08 - 12, 2025

Editors

**Harphool Kumawat, Vineet Kumar, Prashant Shukla,
Aradhana Shrivastava**

Sponsored by

**Board of Research in Nuclear Sciences,
Department of Atomic Energy, Government of India**

National Advisory Committee

R. S. Bhalerao	IISER-PUNE	A. K. Mohanty	DAE
G. Bhattacharyya	SINP	B. K. Nayak	HBNI
S. Bhattacharya	VECC	A. C. Pandey	IUAC
Jayaram N. Chengalur	TIFR	R. G. Pillay	TIFR
A. Chatterjee	BARC	V. S. Ramamurthy	NIAS
R. K. Choudhury	BARC	A. Roy	IUAC
V. M. Datar	IMSc	T. Sakuntala	BARC
A. K. Jain	Amity Univ.	A. Saxena	BARC
B. K. Kanaujia	NITJ	S. Som	VECC
S. Kailas	CEBS	A. K. Sinha	UGC-DAE-CSR
D. Kanjilal	IUAC	D. K. Srivastava	NIAS
S. S. Kapoor	BARC	P. C. Vinod Kumar	SPU, Anand

National Organizing Committee

M. S. Athar	AMU	I. Mazumdar	TIFR
B. R. Behera	PU	H. M. Mittal	NITJ
A. Bharti	UoJ	B. Mohanty	NISER
A. Bhattacharyya	CU	A. Mukherjee	SINP
S. Bhattacharyya	VECC	M. M. Musthafa	Calicut Univ.
R. Chatterjee	IIT-Roorkee	V. Nanal	TIFR
H. Dahiya	NITJ	B. K. Nandi	IITB
J. J. Das	Cotton Univ.	R. Palit	TIFR
P. Das	VECC	A. K. Rai	SVNIT Surat
U. Datta	SINP	R. Sahoo	IIT-Indore
D. Dutta	BARC	S. Santra	BARC
S. S. Ghugre	UGC-DAE CSR	S. Sarkar	VECC
J. A. Gore	BARC	M. K. Sharma	TU Patiala
A. K. Gupta	BARC	A. Shrivastava	BARC
V. Jha	BARC	P. Shukla	BARC
A. Jhingan	IUAC	B. P. Singh	AMU
K. Kalita	GU	R. P. Singh	IUAC
Vineet Kumar	BARC	V. Singh	CUSB
H. Kumawat	BARC	S. K. Tandel	SNIOE
B. Lalremruata	Mizoram Univ.	R. G. Thomas	BARC
K. Mahata	BARC	A. K. Tyagi	BHU
S. K. Mandal	DU		

- A71. Study of multi-quasiparticle band in ^{125}I
Ananya Das, T. Bhattacharjee, A. Saha, S. S Alam, S. Basak, D. Banerjee, A. Bisoi, A. Adhikari, Sangeeta Das, A. Gupta, Y. Sapkota, S. Sharma, D. Pramanik
- A72. Population strength measurements of Se-isotopes using (n_{th}, γ) probe at DURGA facility
A. K. Mondal, A. Chakraborty, P. K. Nayak, U. S. Ghosh, B. Mukherjee, S. Mukhopadhyay
- A73. Reinvestigation of the Lifetime of the Isomeric State in ^{75}Br
Annu Choudhary, Vinod Kumar, Yogendra Pratap Singh, Gobind Ram, Samant Kumar Yadav, Manoj Kumar Sharma, Amritanshu Shukla, Abhishek Yadav, Dhananjaya Sahoo, Ajay Yashavant Deo, Dinesh Negi, Rudrajyoti Palit, Yashraj Jangid, Uday Shankar Ghosh, Indu Bala, Rajesh Pratap Singh
- A74. Measurement of weak branching ratio of ^{159}Dy with ultra-low Q- value
Arindam Sikdar, A. Banerjee, J. Nandi, A. Ray, Parnika Das, T. Bhattacharjee, J. Datta, Devesh Kumar, A. Pal, M. Barman, S. Basak
- A75. Theoretical description of nuclear structure properties of ^{109}Te
Amit Kumar, Rajinder Kumar, Suram Singh, Arun Bharti
- A76. Observation of strong octupole correlations in ^{103}Pd
A. Karmakar, M. Twisha, Ananya Kundu, Piku Dey, Vishal Malik, P. Datta, Satya Samiran Nayak, Saikat Chakraborty, Rudrajyoti Palit, Gopal Mukherjee, Rajarshi Raut, Sanjoy Pal, Sukalyan Chattopadhyay, Sandeep Ghugre
- A77. Investigation of the role of neutron-proton coupling and the search for high spin states in ^{206}Rn
Ajay Kumar, Deepika Choudhury, Sahab Singh, Rajat Roy, Bhoomika Maheshwari, Khamosh Yadav, Rudrajyoti Palit, Vishal Malik, Aditi Sindhu, Shouvik Pal, Shrikant Thorat, Piku Dey, Santosh Telagasetti, Sangeeta Dhuri, Shital Kumar Jadhav, Balaji Naidu, Abraham Thomas, Somnath Nag, Ajay Kumar Singh
- A78. Semidecoupled and vibrational bands in ^{194}Pt
I. Ahmed, S. K. Tandel
- A79. High spin spectroscopy of ^{129}Xe
Anindita Karmakar, Vishal Malik, Rudrajyoti Palit, Saikat Chakraborty, Ananya Kundu, Abraham Thomas, Biswajit Das, Balaji Naidu, Chandrani Majumder, Gopal Mukherjee, Mamta Prajapati, Piku Dey, Pramod Nayak, Sarmishtha Bhattacharyya, Sutanu Bhattacharya, Sneha Das, Shital Kumar Jadhav, Somnath Mukhopadhyay, Satya Samiran Nayak, Shrikant Thorat
- A80. Spectroscopic study of the non-yrast levels in ^{52}Cr
Arkadip Bera, Abhijit Bisoi, R Rahanman, S Sapkota, A Gupta, S Das, S Chakraborty, D Pramanik, S Rajbanshi, A Das, A Pal, S Basak, S Basu, S S Nayek, D Kumar, S Bhattacharya, T Bhattacharjee, G Mukherjee, S Bhattacharyya, A Sharma, S Kundu, P Giri, R Raut, S S Ghugre

Spectroscopic study of the non-yrast levels in ^{52}Cr

Arkadip Bera,¹ Abhijit Bisoi,^{1,*} Rozina Rahaman,¹ Y. Sapkota,² Arkabrata Gupta,³
 Sangeeta Das,³ S. Chakraborty,³ D. Pramanik,⁴ S. Rajbanshi,⁵ Ananya Das,^{6,7}
 A. Pal,^{6,7} S. Basak,^{6,7} Sansaptak Basu,^{6,7} S. S. Nayak,^{6,7} D. Kumar,^{6,7}
 Soumik Bhattacharya,^{6,7} T. Bhattacharjee,^{6,7} G. Mukherjee,^{6,7} S. Bhattacharyya,^{6,7}
 A. Sharma,⁸ S. Kundu,⁸ Pankaj K. Giri,⁸ R. Raut,⁸ and S. S. Ghugre⁸

¹Indian Institute of Engineering Science and Technology, Shibpur, Howrah 711103, India

²Department of Physics, Dudhnoi College, Dudhnoi, Goalpara, Assam 783124, India

³Institute of Engineering and Management, UEM, Kolkata, Kolkata 700160, India

⁴Haldia Institute of Technology, Haldia 721657, India

⁵Presidency University, Kolkata 700073, India

⁶Variable Energy Cyclotron Centre, 1/AF Bidhannagar, Kolkata 700064, India

⁷Homi Bhabha National Institute, Training School Complex, Anushaktinagar, Mumbai 400094, India

⁸UGC-DAE Consortium for Scientific Research, Kolkata Centre, Kolkata 700098, India

*Email: abhijitbisoi@physics.iests.ac.in

Introduction

The spectroscopic study of $1f_{7/2}$ shell nuclei near $N \sim 28$ is of significant interest because these nuclei serve as ideal candidates for description within the shell model framework. They exhibit a diverse range of nuclear phenomena, including collective behavior, band termination, and spherical-deformed shape coexistence. These phenomena have been effectively explained using advanced γ -ray spectroscopy techniques and have shown good agreement with shell model calculations.

The nucleus ^{52}Cr possesses 12 valence nucleons (4 protons and 8 neutrons) in the $1f_{7/2}$ shell. Numerous studies have been conducted on this nucleus using both heavy-ion and light-ion reactions. The most recent work on ^{52}Cr , populated through heavy ion induced reaction, was reported by Rajesh Kumar *et al.* [1]. In this study, the level scheme was extended up to 10 MeV, with spins reaching $J^\pi = 13^+$ and investigated only the structural properties of the yrast levels of ^{52}Cr . The primary objective of the present study is to extend the level scheme of ^{52}Cr horizontally and investigate the structural properties of the non-yrast levels in ^{52}Cr . Finally, the experimental observation will be compared with the results obtained from shell model calculations to understand their microscopic origin.

Experimental Details

High-spin states of ^{52}Cr were populated via $^{51}\text{V}(^4\text{He}, 2n)^{52}\text{Cr}$ fusion evaporation reaction.

The experiment was performed using a 45 MeV alpha beam, delivered from the K-130 Cyclotron, stationed at the Variable Energy Cyclotron Centre (VECC), Kolkata. A self-supported $\sim 12 \text{ mg/cm}^2$ thick natural vanadium target was used for the experiment, within which $\sim 95\%$ of the recoils stopped. The de-excitation of γ -rays was detected using the Indian National Gamma Array (INGA) at VECC, which was comprised of 11 Compton-suppressed clovers (6 were mounted at 90° , 3 at 125° , and 2 at 40°) and 1 LEPS (mounted at 40°) at the time of experiment. Singles and two-fold coincidence events were recorded in list mode using a PIXIE-16 based digital data acquisition system [2]. The raw data were sorted, and both angle-independent and angle-dependent symmetric and asymmetric matrices were constructed using the BiNDAS package [3]. Data analysis was carried out using INGASORT [4]. Energy calibration was performed with the radioactive source ^{152}Eu and some online gammas having energies up to 3 MeV. The relative efficiency of the clover detectors was determined using radioactive sources ^{152}Eu and ^{66}Ga .

Results and Discussions

The excited states of ^{52}Cr were investigated through coincidence relationships and relative intensity measurements using a symmetric 90° vs 90° matrix, with careful consideration given to the angular distribution of dipole and quadrupole transitions. All existing transitions were observed by putting a gate on the 1433.9 keV ground state transition (Fig. 1). We have not

found any transitions coming from the levels above $J^\pi=13^+$. But several new transitions have been found, which are mostly coming from the non-yrast levels. We have placed 19 new transitions and 8 new levels in the level scheme based on their coincidence relationship and relative intensities.




Fig. 1: Typical background-subtracted spectrum [(a) 300 to 1730 keV, (b) 1730 to 4096 keV] obtained by putting a gate on the 1433.9 keV transition. It shows all the existing (black) and new (red) transitions that belong to ^{52}Cr . The spectrum was generated from 90° vs. 90° matrix.

The spin-parity of most of the newly identified levels has been confirmed based on the results obtained from Directional correlation (DCO) and polarization (IPDCO) measurements. The DCO and IPDCO measurements could not be performed for a few new transitions due to their poor statistics. The spin and parity of a few levels were earlier assigned tentatively [1]. In the present work, we have confirmed / modified the spin and parity of those levels using their measured DCO and IPDCO values of the decay out transitions of interest.

We have observed Doppler shifted peaks for a few transitions in ^{52}Cr , as shown in Fig. 2. Lineshape analysis will therefore be carried out to measure the lifetimes of these levels.




Fig. 2: Doppler shifted peaks of (a) 1035.1 and (b) 1614.8 keV transitions at three different angles.

Large basis shell model calculations were performed using the NUSHELLX code [5] and KB3GPN [6] effective interaction. The valence space consisted of the full fp major shell with $1f_{7/2}$, $1f_{5/2}$, $2p_{3/2}$, and $2p_{1/2}$ orbitals for both protons and neutrons above the ^{40}Ca inert core. The calculations were performed without any particle restrictions in the fp valence space. The calculated energies have been compared with the experimental energies and they are mostly agreed well. Now, transition probabilities, moments, spectroscopic factors, etc., will be carried out to understand the microscopic structure of the non-yrast levels of ^{52}Cr .

Acknowledgements

The authors sincerely thank the accelerator staff of VECC, Kolkata for a nearly uninterrupted beam and also thank the INGA (partially funded by the Department of Science and Technology, Government of India) collaboration.

References

- [1] Rajesh Kumar *et al.*, Phys. Rev. C **76**, 034301 (2007).
- [2] S. Das *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **893**, 138 (2018).
- [3] S.S. Nayak, G. Mukherjee IEEE Trans. Nucl. Sci., **70** (2023), p. 2561
- [4] R. Bhowmik *et al.*, in Proc. DAE Symp. Nucl. Phys., Vol. 44B (2001) p. 422.
- [5] W. D. M. Rae, NUSHELLX, <http://www.garsington.eclipse.co.uk>.
- [6] A. Poves, J. Sánchez-Solano, E. Caurier, and F. Nowacki, Nucl. Phys. A **694**, 157 (2001).



ISBN 978-81-984360-5-4



9 788198 436054